

STARMEC[®] LV POUR-ON FOR CATTLE

STARMEC[®] LV POUR-ON FOR CATTLE contains Ivermectin, which has a broad spectrum of activity against most internal and external parasites with sensitivity to Ivermectin. Ivermectin is a member of the macrocyclic lactones (anthelmintic family of chemicals). Resistance may develop to any chemical.

Indications: For the treatment and control of adult and some immature stages of:

Gastro-intestinal roundworms

Barber's Pole Worm (*Haemonchus placei*)
Hookworm (*Bunostomum phlebotomum*)
Intestinal Threadworm (*Strongyloides papillosus*)
Large Bowel Worm (*Oesophagostomum radiatum*)
Small Intestinal Worm (*Cooperia oncophora*)
(*Cooperia punctata*)

Small Brown Stomach Worm (*Ostertagia ostertagi*)
(including inhibited stages)

Stomach Hair Worm (*Trichostrongylus axei*)

Thin Necked Intestinal Worm

(*Nematodirus spathiger*)

Whipworm (*Trichuris* spp.)

Lungworms (*Dictyocaulus viviparus*)

Sucking and Biting Lice

Biting lice (*Bovicola bovis*)

Little sucking lice (*Solenopotes capillatus*)

Long nosed sucking lice (*Linognathus vituli*)

Short nosed sucking lice

(*Haematopinus eurysternus*)

Mites (*Chorioptes bovis*, *Sarcoptes scabiei var bovis*)

Buffalo Fly (*Hoematobia irritans exigua*)

Cattle Tick (*Boophilus microplus*)

Maximum tick efficacy is achieved 4-5 days after treatment. In planning a strategic tick control program, it should be noted that viable eggs may have been laid by engorged female ticks that drop from cattle in the initial days after treatment.

STARMEC[®] LV POUR-ON FOR CATTLE should be alternated with conventional dips or Pour-Ons when the program requires repeat treatments.

Sustained Activity:

STARMEC[®] LV POUR-ON FOR CATTLE applied at the recommended dose rate of 1mL per 20kg bodyweight effectively controls infection with *Ostertagia* spp. (Small brown stomach worm) acquired up to 14 days after treatment, with *Dictyocaulus viviparus* (Lungworm) acquired up to 28 days after treatment and *Haematobia irritans exigua* (Buffalo Fly) up to 21 days after treatment.

DIRECTIONS FOR USE:

Dosage and Administration:

STARMEC[®] LV POUR-ON FOR CATTLE is for external application to cattle only.

The recommended dose rate for STARMEC[®] LV POUR-ON FOR CATTLE is 1 mL/20 kg body weight.

Apply along the topline of the animal.

Cattle heavier than 650 kg should be dosed at 1 mL per 20 kg.

A representative sample of animals should be weighed before treatment either with scales or a weighband.

Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves, etc.). Do not under dose.

Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

Liveweight (kg)	Dose (mL)	No head treated per pack
51 – 100	5.0	2000
101 – 150	7.5	1333
151 – 200	10.0	1000
201 – 250	12.5	800
251 – 300	15.0	666
301 – 350	17.5	571
351 – 400	20.0	500
401 – 450	22.5	444
451 – 500	25.0	400
501 – 550	27.5	363
551 – 600	30.0	333
601 – 650	32.5	307

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER THE APPROPRIATE LEGISLATION.



STARMEC[®] LV POUR-ON FOR CATTLE

ACTIVE CONSTITUENT: 10g/L IVERMECTIN



For the treatment and control of ivermectin sensitive internal and external parasites of beef and dairy cattle.

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

General directions

Rainfall on the herd within 2 hours of application may result in reduced efficacy of the product. Cattle should not be treated with STARMEC[®] LV POUR-ON FOR CATTLE when the hide or hair is wet since the efficacy of the product may be reduced. Temporary hair loss occasionally occurs in the application area but will resolve without further treatment.

Do not retreat with STARMEC[®] LV POUR-ON FOR CATTLE within the meat withholding period of 42 days.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 42 days before slaughter for human consumption.

MILK: Zero (0) days.

TRADE ADVICE - EXPORT SLAUGHTER INTERVAL (ESI): DO NOT use less than 42 days before slaughter for export. Before using this product, confirm the current ESI from The Hunter River Company on (02) 9499 7370 or the APVMA website (www.apvma.gov.au/residues).

SAFETY DIRECTIONS: Poisonous if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and goggles. If product gets into the eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

ADDITIONAL USER SAFETY INFORMATION: Additional information is listed in the safety data sheet.

ENVIRONMENTAL PROTECTION: Ivermectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL: Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Store below 30°C (Room Temperature).

APVMA Approval No.: 83891/109193

Batch:

Expiry:



The Hunter River Company Pty Limited

7/20 St Johns Avenue,
Gordon NSW, 2072.

Ph: (02) 9499 7370

www.hunterriverco.com.au

STARMEC is a trademark of
The Hunter River Company Pty Limited



THE HUNTER RIVER CO.

File Name	StarmecLV Pour-on_10L.pdf
Date	08.02.2018
Full-Size Printed Label Dimensions	400 x 100 mm
Approximate Scale (%) at which the pdf will print on A4 (or A3)	A4 : 57% A3 : 85%