



Company name:	THE HUNTER RIVER COMPANY PTY LIMITED
Product name:	STARMEC LV POUR-ON FOR CATTLE
e-Label application No:	N/A
APVMA Approval No:	83891 / 114051

Product No:	83891	Version No:	1.0
Application started:		Version Created:	
Started By:		Printed:	

Label name:	STARMEC LV POUR-ON FOR CATTLE
-------------	-------------------------------

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	---

Constituent Statements:	ACTIVE CONSTITUENT: 10 g/L Ivermectin Also contains: 151 g/L Diethylene glycol monobutyl ether (solvent)
-------------------------	---

Claims:	<p>For the treatment and control of Ivermectin sensitive internal and external parasites of beef and dairy cattle.</p> <p>STARMEC LV POUR-ON FOR CATTLE contains Ivermectin, which has a broad spectrum activity of against most internal and external parasites with sensitivity to Ivermectin. Ivermectin is a member of the macrocyclic lactones (anthelmintic family of chemicals). Resistance may develop to any chemical.</p> <p>Indications: For the treatment and control of adult and some immature stages of: Gastro-intestinal roundworms Barber's Pole Worm (<i>Haemonchus placei</i>) Hookworm (<i>Bunostomum phlebotomum</i>) Intestinal Threadworm (<i>Strongyloides papillosus</i>) Large Bowel Worm (<i>Oesophagostomum radiatum</i>) Small Intestinal Worm (<i>Cooperia oncophora</i>) (<i>Cooperia punctata</i>) Small Brown Stomach Worm (<i>Ostertagia ostertagi</i>) (Including inhibited stages) Stomach Hair Worm (<i>Trichostrongylus axei</i>) Thin Necked Intestinal Worm (<i>Nematodirus spathiger</i>) Whipworm (<i>Trichuris</i> spp.)</p> <p>Lungworms (<i>Dictyocaulus viviparus</i>)</p> <p>Sucking and Biting Lice Biting lice (<i>Bovicola bovis</i>) Little sucking lice (<i>Solenopotes capillatus</i>) Long nosed sucking lice (<i>Linognathus vituli</i>)</p>
---------	--

	<p>Short nosed sucking lice (<i>Haematopinus eurysternus</i>) Mites (<i>Chorioptes bovis</i>) (<i>Sarcoptes scabiei</i> var <i>bovis</i>)</p> <p>Buffalo Fly (<i>Haematobia irritans exigua</i>)</p> <p>Cattle Tick (<i>Rhipicephalus [Boophilus] microplus</i>)</p> <p>Maximum tick efficacy is achieved 4-5 days after treatment. In planning a strategic tick control program, it should be noted that viable eggs may have been laid by engorged female ticks that drop from cattle in the initial days after treatment.</p> <p>STARMEC LV POUR-ON FOR CATTLE should be alternated with conventional dips or Pour-Ons when the program requires repeat treatments.</p> <p>Sustained Activity: STARMEC LV POUR-ON FOR CATTLE applied at the recommended dose of 1 mL per 20 kg bodyweight effectively controls infection with <i>Ostertagia</i> spp. (Small brown stomach worm) acquired up to 14 days after treatment, with <i>Dictyocaulus viviparus</i> (Lungworm) acquired up to 28 days after treatment and <i>Haematobia irritans exigua</i> (Buffalo Fly) up to 28 days after treatment.</p>
Net Contents:	500 mL, 2.5 L, 10 L, 12.5 L, 15 L, 20 L, 25 L
Directions for Use:	
Restrains:	Re-treatment interval: DO NOT re-treat animals for 42 days after last treatment.
Contraindications:	
Precautions:	
Side effects:	
Dosage and Administration:	<p>STARMEC LV POUR-ON FOR CATTLE is for external application to cattle only.</p> <p>The recommended dose rate for STARMEC LV POUR-ON FOR CATTLE is 1 mL/20 kg body weight. Apply along the topline of the animal.</p> <p>Liveweight (kg); Dose Volume (mL)</p> <p>51-100; 5.0 101-150; 7.5 151-200; 10 201-250; 12.5 251-300; 15.0 301-350; 17.5 351-400; 20.0 401-450; 22.5 451-500; 25.0</p>

	<p>501-550; 27.5 551-600; 30.0 601-650; 32.5</p> <p>Cattle heavier than 650 kg should be dosed at 1 mL per 20 kg. A representative sample of animals should be weighed before treatment either with scales or a weighband.</p> <p>Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves, etc.). Do not under dose.</p> <p>Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.</p>
General Directions:	<p>NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.</p> <p>Rainfall on the herd within 2 hours of application may result in reduced efficacy of the product. Cattle should not be treated with STARMEC LV POUR-ON FOR CATTLE when the hide or hair is wet since the efficacy of the product may be reduced. Temporary hair loss occasionally occurs in the application area but will resolve without further treatment.</p>
Withholding Periods:	<p>MEAT: DO NOT USE less than 42 days before slaughter for human consumption. MILK: Zero (0) days.</p>
Trade Advice:	<p>EXPORT SLAUGHTER INTERVAL (ESI): DO NOT use less than 42 days before slaughter for export. Before using this product, confirm the current ESI from The Hunter River Company on (02) 9499 7370 or the APVMA website (www.apvma.gov.au/residues).</p>
Safety Directions:	<p>Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.</p>
First Aid Instructions:	<p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.</p>
First Aid Warnings:	
Additional User Safety:	<p>Additional information is listed in the safety data sheet.</p>
Environmental Statements:	<p>ENVIRONMENTAL PROTECTION Ivermectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.</p>
Disposal:	<p>[500mL] Dispose of container by wrapping with paper and putting in garbage.</p>

	<p>[2.5L, 10L, 12.5L, 15L, 20L, 25L] Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p>
--	---

Storage:	Store below 30°C (room temperature).
----------	--------------------------------------