

**POISON**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



**Dow AgroSciences**

# Grazon™ Extra

## Herbicide

**ACTIVE CONSTITUENTS:** 300 g/L TRICLOPYR present as butoxyethyl ester  
100 g/L PICLORAM present as hexyloxypropylamine salt  
8 g/L AMINOPYRALID present as hexyloxypropylamine salt

**GROUP I HERBICIDE**

For control of a range of environmental and noxious woody and herbaceous weeds as specified in the Directions For Use table.

**SHAKE WELL BEFORE USE.**

**Dow AgroSciences Australia Limited** ABN 24 003 771 659

20 Rodborough Road FRENCHS FOREST NSW 2086

[www.dowagrosciences.com.au](http://www.dowagrosciences.com.au)

**CUSTOMER SERVICE TOLL FREE 1-800 700 096**

®™ Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow



**Dow AgroSciences**

## DIRECTIONS FOR USE

### RESTRAINTS

- **DO NOT** apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition, presence of disease, damage or previous herbicide treatment, as reduced levels of control may result.
- **DO NOT** spray if rain is likely within 1 hour or if foliage is wet from rain or dew.
- **DO NOT** burn off, cut or clear blackberry or other woody weeds for at least 6 months after spraying.
- **DO NOT** apply by aerial application in wind in excess of 15 km/hr and/or air temperatures above 35°C.
- In areas prone to flooding treatment should commence after any annual flooding as such areas flooded within 9 months following application may have reduced results.
- **DO NOT** make hay or silage, from pastures treated with Grazon Extra, for 6 months after application.

### 1. WOODY WEED SITUATIONS

**Table A: High Volume Spraying**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

**Note:** Dow AgroSciences only recommends the use of knapsacks or 12 volt sprayers for use on low growing herbaceous weeds and woody weeds that are not regrowth less than 60 cm high or 60 cm diameter.

| AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY. |   |                        |                   |   |
|--|---|------------------------|-------------------|---|
| WEEDS CONTROLLED   | WEED GROWTH STAGE                           | STATE                  | RATE /100 L Water | CRITICAL COMMENTS   |
| African boxthorn   | Less than 2 m tall                          | All States             | 500 mL            | Apply when bushes have good leaf cover, growth and <b>no leaf fall</b> .  |
| <i>Angophora</i> spp.  | 1 to 3 m tall                               |                        | 350 mL            |   |
| Australian blackthorn  | Less than 2 m tall                          |                        | 500 mL            | Apply from late spring to early autumn.   |
| <i>Banksia</i> spp.  | 1 to 3 m tall                               |                        | 350 mL            |   |
| Biddy bush (Chinese shrub) (Sifton bush)   | Autumn when actively growing                | ACT, NSW only          | 500 mL            | Add BS-1000 or an alternative (see compatibility section) at 125 mL/100 L of water for best results.  |
| Blackberry in association with:<br>Docks<br>Ragwort<br>Smartweed<br>Thistles                       | Late spring to autumn                       | All States             | 350 mL or 500 mL  | Use the higher rate on plants that have been damaged by grazing stock or insects and on known difficult to kill blackberry.<br><br>Where herbicides other than Group I herbicides have been used, allow 2 seasons regrowth to occur before respraying with Grazon™ Extra. |
| Blue heliotrope  | Flowering                                   | ACT, NSW, NT, Qld only | 500 mL            | Apply in a minimum spray volume of 1250 L/ha.   |
| Brooms:<br>Cape English<br>Flax leaf<br>Montpellier  | Spring to mid-summer prior to pod formation | All States             | 250 mL            | Apply as a thorough foliage spray.  |
|  | Autumn to winter                            |                        | 350 mL            |   |
| Camphor Laurel   | Less than 2 m tall                          | All States             |                   |   |
|  | Above 2 m tall                              |                        | 500 mL            |   |
| Capeweed   | Flowering                                   |                        | 150 mL            |   |
| <i>Casuarina</i> spp.  | 1 to 3 m tall                               |                        | 350 mL            |   |



**Table A: High Volume Spraying** *continued*

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

| AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY. |   |                        |                   |  |                                    |
|--|---|------------------------|-------------------|--|------------------------------------|
| WEEDS CONTROLLED   | WEED GROWTH STAGE   | STATE                  | RATE /100 L Water | CRITICAL COMMENTS  |                                    |
| Chinee apple   | Less than 2 m tall  | NT, Qld, WA only       | 350 mL            | Add BS-1000 or an alternative (see compatibility section) at 100 mL/100 L of water for best results.   |                                    |
| Cockspur thorn   | Spring to autumn  | ACT, NSW, NT, Qld only |                   | Apply as a thorough foliage spray.   |                                    |
| Common sensitive plant   | Any time when actively growing                              | NT, Qld, WA only       | 200 mL            | To avoid leaves closing during application, spray plants while moving forward. Add BS-1000 or an alternative (see compatibility section) at 100 mL/100 L of water for best results.        |                                    |
| Crofton weed   | Spring to autumn  | ACT, NSW, NT, Qld only | 350 mL            | Apply as a thorough foliage spray.   |                                    |
| Eastern cotton bush ( <i>Maireana microphylla</i> )  | Spring to autumn  | ACT, NSW NT, Qld only  | 500 mL            | Add Uptake™ Spraying Oil at 500 mL/100 L of water. Some bushes may require a follow-up spray to control regrowth.  |                                    |
| <i>Eucalyptus</i> spp.   | Seedling and regrowth from small lignotubers, 1 to 3 m tall | All States             | 350 mL or 500 mL  | <b>Apply the high rate where difficult to control species of <i>Eucalyptus</i> regrowth is present.</b> Addition of an adjuvant may improve results. Contact Dow AgroSciences for details. |                                    |
| Fireweed   | Flowering   |                        | 350 mL            |  | Apply as a thorough foliage spray. |
| Galenia  | Fresh growth in spring to summer                            |                        | 500 mL            |  | Use 2000 L of spray mixture/ha.    |
| Giant bramble  | Spring to autumn  | NT, Qld, WA only       |                   | Penetration of thick clumps may be difficult and respraying may be necessary. Add BS-1000 or an alternative (see compatibility section) of water for best results.                         |                                    |
| Gorse  | 1 to 1.5 m tall   | All States             | 250 mL            | Spring and summer treatment only. Add BS-1000 or an alternative (see compatibility section) at 100 mL/100 L of water for best results.   |                                    |
|  | Over 1.5 m tall or autumn treatment                         |                        | 350 mL            | Add BS-1000 or an alternative (see compatibility section) at 100 mL/100 L of water for best results.   |                                    |
|  | Winter treatment  |                        | 500 mL            | Brownout may not be complete until summer. Add BS-1000 or an alternative (see compatibility section) at 100 mL/100 L of water for best results.  |                                    |
| Groundsel bush ( <i>Baccharis halimifolia</i> )  | 1 to 1.5 m tall in spring to summer                         | All States             | 250 mL            | Apply as a thorough foliar spray.  |                                    |
|  | Over 1.5 m tall or autumn treatment                         |                        | 350 mL            |  |                                    |
| Green cestrum  | Late spring to early autumn                                 | ACT, NSW, NT, Qld only | 500 mL            | One application may give satisfactory control. Any subsequent regrowth and seedlings must be resprayed at approximately 1 metre high.  |                                    |
| Hawthorn   | Less than 2 m tall  | All States             |                   | Apply from late spring to early autumn.  |                                    |



**Table A: High Volume Spraying** *continued*

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

| AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY. |   |                            |                   |   |
|--|---|----------------------------|-------------------|---|
| WEEDS CONTROLLED   | WEED GROWTH STAGE                                       | STATE                      | RATE /100 L Water | CRITICAL COMMENTS   |
| Horehound  | Pre-flowering   | All States                 | 350 mL            | Apply as a thorough foliar spray.   |
| Japanese sunflower   |   | ACT, NSW, NT, Qld only     |                   |   |
| Lantana<br>( <i>Lantana camara</i> )<br>( <i>Lantana montevidensis</i> )                           | Up to 1 m tall in summer to autumn                      | All States                 | 500 or 750 mL     | Add one of the following adjuvants, when using 350 mL rate:<br>Uptake Spraying Oil @ 0.5% v/v.<br>Pulse® Penetrant @ 0.1% v/v.  |
|  | 1 to 2 m tall in summer to autumn                       |                            |                   | Thoroughly wet foliage stems and soil around the base of the plants. Use higher rate on known harder to kill varieties.   |
| Lion's tail<br>( <i>Leonatis nepetifolia</i> )   | Pre-flowering   | NT, Qld only               | 200 mL            | Apply as a thorough foliar spray.<br>Add BS-1000 or an alternative (see compatibility section) at 100 mL/100 L of water for best results.                             |
| Limebush   | Any time of year with good leaf cover and soil moisture | NT, Qld only               | 350 mL            | Penetration of thick clumps may be difficult and respraying may be required.<br>Addition of an adjuvant may improve results.<br>Contact Dow AgroSciences for details. |
| Manuka   | At flowering  | Vic only                   | 500 mL            | For optimum results, add Pulse Penetrant at 200 mL/100 L of spray.<br>Thoroughly wet foliage, stems and soil around the base of the plants.                           |
| Mesquite<br>( <i>Prosopis</i> spp.)  | Seedling, full leaf and flowering before podding        | ACT, NSW, NT, Qld, WA only | 350 mL            | <b>DO NOT</b> spray plants bearing pods.<br>Add BS-1000 or an alternative (see compatibility section) at 100 mL/100 L of water for best results.                      |
| <i>Prosopis velutina</i>   |   | NT, Qld only               | 670 mL            |   |
| Milfoil (Yarrow)   | Flowering   | ACT, NSW only              | 350-500 mL        | Use low rate when in close proximity to highly sensitive vegetation.  |
| Mistflower   | Spring to Autumn  | ACT, NSW, NT, Qld only     | 350 mL            | Apply as a thorough foliar spray.   |
| Mother-of-millions   | Flowering   | ACT, NSW, NT, Qld only     | 500 mL            | Add BS-1000 or an alternative (see compatibility section) at 100 mL/100 L of water.   |
| Paddy's lucerne  | Active growth   | ACT, NSW only              |                   | Plants that have been continually slashed or grazed over many seasons may be difficult to control and regrowth may occur.   |
| Parkinsonia  | Under 2 m tall  | NT, Qld, WA only           | 350 mL            | Add Uptake Spraying Oil at 500 mL/100 L water.<br>Avoid spraying under dry conditions when plants are stressed or bearing pods. Thoroughly wet foliage.               |



**Table A: High Volume Spraying** *continued*

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

| AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY. |   |                        |                   |  |
|--|---|------------------------|-------------------|--|
| WEEDS CONTROLLED   | WEED GROWTH STAGE                                 | STATE                  | RATE /100 L Water | CRITICAL COMMENTS  |
| Paterson's curse   | Flowering   | All States             | 250 mL            |  |
| Prickly pear (common)<br>Smooth tree pear  | Active phyllode growth                            |                        | 500 mL            | Apply as a thorough foliage spray. Regrowth may occur, so a follow-up application may be necessary.  |
| Rubber vine (Not infected with rust)   | Up to 1.5 m tall at flowering                     | NT, Qld, WA only       | 350 mL            | Spray all leaves and stems just to the point of runoff and thoroughly spray the base of the plant. With larger, more dense stands, regrowth may occur. Subsequent control of any regrowth should be done by basal bark spraying. |
|  | Dense stands greater than 1.5 m tall at flowering |                        | 500 mL            |  |
| Siam weed  | Active growth                                     | NT, Qld, WA only       | 350 mL            | Add BS-1000 or an alternative (see compatibility section) at the rate of 100 mL/100 L water for best results.  |
| Sicklepod  | Up to flowering                                   | NT, Qld, WA only       | 200 mL            | <b>DO NOT</b> apply to podding plants. Add BS-1000 or an alternative (see compatibility section) at the rate of 100 mL/100 L water for best results.   |
| Spear thistle  | Rosette to flowering                              | All States             | 150 mL            | Late spring to early summer.   |
| St John's wort   | From flowering to early seed set                  |                        | 500 mL            |  |
| Sweet briar  | Up to 1.5 m tall                                  |                        | 350 mL            | Add metsulfuron-methyl at 10g/100 L water to obtain more reliable results with the lower rate of Grazon Extra. Full leaf to ripe fruit prior to leaf fall. Thorough wetting including the crown is recommended.                  |
|  |   |                        | 500 mL            |  |
| Tobacco weed   | Actively growing plants                           | NT, Qld, WA only       | 300 mL            | Add BS-1000 or an alternative (see compatibility section) at 100 mL/100 L of water for best results.   |
| Wattle ( <i>Acacia</i> spp.) (except corkwood wattle)  | 1 to 3 m tall                                     | All States             | 350 mL            |  |
| Wild Rosemary ( <i>Cassinia laevis</i> )   | Active growth 0.5 to 1.0 m tall                   | NT, Qld only           | 350 to 500 mL     | Use lower rate on seedlings 0.5 m tall. Apply as a thorough foliar spray.  |
| Wild tobacco tree  | Spring to autumn up to 2 m tall                   | ACT, NSW, NT, Qld only | 350 mL            |  |



**Table B: Aerial Application**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

| <b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, PASTURES AND RIGHTS-OF-WAY.</b> |   |  |   |  |
|--|---|--|---|--|
| <b>WEEDS CONTROLLED</b>  | <b>WEED GROWTH STAGE</b>                  | <b>STATE</b>                             | <b>RATE /ha</b>                                   | <b>CRITICAL COMMENTS</b>   |
| Blackberry   | Summer to autumn                          | ACT, NSW, NT, Qld, SA, Vic, WA only      | 10 L  | Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before respraying with Grazon Extra.<br><b>WARNING:</b> <i>Eucalyptus</i> species up to 4 m may be killed if sprayed during this treatment. Mature trees which are 15 to 20 m tall may be partially defoliated but are likely to recover. |
| Goose  |   | Tas only                                 |   | Helicopter application only.   |
| Cockspur thorn<br>Crofton weed<br>Lantana<br>Mistflower  | Late autumn                               | ACT, NSW, NT, Qld only (helicopter only) | 1.5 L + 6.L 2,4-D 625 amine (625 g/L formulation) | Spray with calibrated equipment using the half overlap opposite pass technique applying a minimum spray volume of 150 L/ha.<br><b>Follow-up respraying will be required.</b>   |
| Lantana  |   |  | 10 L  | Helicopter application only.   |
| Rubber vine (Not infected with rust)   | When flowering                            | NT, Qld only (helicopter only)           | 3 L to 5 L  | Use rates will depend upon the density and height of the rubber vine stand. The higher rate should be used on dense stands, however, complete coverage and penetration may be difficult.<br><b>Follow-up respraying will be required.</b><br><br>Any regrowth should be sprayed with a suitable basal bark herbicide.                        |
| St John's wort   | Flowering to early seed set (Nov-Jan)     | ACT, NSW only                            | 4 L   | Helicopter application only. Follow-up spraying will be required in the following season.  |
| <b>AGRICULTURAL NON-CROP AREAS ON FLOOD PLAINS.</b>  |   |  |   |  |
| <b>WEEDS CONTROLLED</b>  | <b>WEED GROWTH STAGE</b>                  | <b>STATE</b>                             | <b>RATE /ha</b>                                   | <b>CRITICAL COMMENTS</b>   |
| Parkinsonia  | Seedlings 1-2 m tall, or 12-24 months old | Qld and NT only (helicopter only)        | 3 L   | Add Uptake Spraying Oil at 1 L/ha.   |

**Table C: Controlled Droplet Application (C.D.A.)**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

| <b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b> |                          |              |                 |  |
|---|--------------------------|--------------|-----------------|--|
| <b>WEEDS CONTROLLED</b>   | <b>WEED GROWTH STAGE</b> | <b>STATE</b> | <b>RATE</b>     | <b>CRITICAL COMMENTS</b>   |
| Blackberry in association with:<br>Docks<br>Ragwort<br>St John's wort<br>Thistles                         | Summer to autumn         | All States   | Apply undiluted | One application may give satisfactory control but subsequent regrowth and seedlings should be resprayed after hardening off. Where herbicides other than Group I herbicides have been used, allow 2 seasons regrowth to occur before respraying with Grazon Extra. |



**Table D: Low Volume High Concentrate Application Techniques (Gas Powered Gun, Sprinkler Sprayer)**  
See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

| <b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b> |                                     |                         |                         |  |
|---|-------------------------------------|-------------------------|-------------------------|--|
| <b>WEEDS CONTROLLED</b>   | <b>WEED GROWTH STAGE</b>            | <b>STATE</b>            | <b>RATE /10 L Water</b> | <b>CRITICAL COMMENTS</b>   |
| Blackberry  | Late spring to autumn               | All States              | 335 mL                  | Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high volume application technique is recommended. |
| Camphor Laurel<br>Cockspur thorn<br>Crofton weed  | Less than 1.5 m high                | ACT, NSW, NT, Qld only  | 500 mL                  |  |
| <i>Eucalyptus</i> species   | Seedlings up to 2 m tall            | All States              | 335 mL                  |  |
| Mistflower  | Less than 1.5 m high                | ACT, NSW, NT, Qld only  | 500 mL                  |  |
| Sweet briar   | 1.5 m tall, full leaf to ripe fruit | ACT, NSW only           |                         | <b>Gas Powered Gun only:</b> Apply to actively growing bushes not more than 1.5 m tall that have not more than 5 stems from the crown.               |
| St John's wort  | During flowering to early seed set  | ACT, NSW, Tas, Vic only |                         | <b>Gas Powered Gun only:</b> One application should provide control. Minor regrowth and seedlings may be retreated the following summer.             |
| Wild tobacco tree   | Less than 1.5 m high                | ACT, NSW, NT, Qld only  |                         | Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high volume application technique is recommended. |

**Table E: Boom Application**  
See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

| <b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b> |                                       |               |                 |  |
|---|---------------------------------------|---------------|-----------------|--|
| <b>WEEDS CONTROLLED</b>   | <b>WEED GROWTH STAGE</b>              | <b>STATE</b>  | <b>RATE /ha</b> | <b>CRITICAL COMMENTS</b>   |
| Galenia   | Fresh growth during spring to summer  | All States    | 5 L             | Rough mine sites will require adequate spray equipment such as boomless nozzles for effective coverage.  |
| Sicklepod   | Up to flowering                       | NT, Qld only  | 3 L             | <b>DO NOT</b> apply to podding plants. Add BS-1000 or an alternative (see compatibility section) at the rate of 100 mL/100 L water.                    |
| St John's wort  | Flowering to early seed set (Nov-Jan) | ACT, NSW only | 2 to 4 L        | Use the higher rate on dense infestations and when longer residual control is required. Follow-up respraying will be required in the following season. |



## 2. FALLOW SITUATIONS

**Table A: Boom Application**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

| FALLOW  |   |                                 |   |   |
|---|---|---------------------------------|---|---|
| WEEDS CONTROLLED  | WEED GROWTH STAGE   | STATE                           | RATE /ha  | CRITICAL COMMENTS   |
| Blackberry<br>nightshade<br>- <b>suppression only</b>                           | 10 to 25 cm tall,<br>prior to flowering                         | ACT,<br>NSW,<br>NT, Qld<br>only | 200 to 400 mL<br>1.125 L + Ripper™ 480<br>(glyphosate) + adjuvant | <p><b>For use by ground equipment only.</b><br/>Plants must be actively growing.</p> <p>Use the lower rate on the smaller weeds, as specified in the weed growth stage (or up to 5 cm diameter for <i>Polymeria pusilla</i>).</p> <p>Refer to Ripper 480 label for use of adjuvant.</p> <p>For Northern New South Wales and Queensland, <b>DO NOT</b> plant susceptible crops for up to 9 months after application, as specified in <b>GENERAL INSTRUCTIONS - MINIMUM RECROPPING PERIODS – Table A.</b></p> <p>Dry conditions after application will increase the re-cropping interval.</p> <p>For Southern New South Wales, please refer to <b>MINIMUM RECROPPING PERIODS – Table C.</b></p> |
| Camel melon<br>Prickly paddy melon<br>Cucumber melon<br>( <i>Cucumis melo</i> ) | From 2 leaf to<br>50 cm diameter                                |                                 |   |   |
| Common sowthistle   | From 8 leaf to<br>flowering                                     |                                 |   |   |
| Cow vine  | From 2 to 5 leaf up<br>to 15 cm diameter,<br>prior to flowering |                                 | 300 to 500 mL + 1.125 L<br>Ripper™ 480 (glyphosate)<br>+ adjuvant |   |
| Lucerne (established)   | Active growth,<br>15 to 25 cm high,<br>during spring            |                                 |   |   |
| <i>Polymeria pusilla</i>  | 2 to 12 leaf up to<br>20 cm diameter,<br>prior to flowering     |                                 |   |   |

**Table B: Blanket Wiper Application**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

| FALLOW                                       |                                 |                 |   |   |
|--|---------------------------------|-----------------|---|---|
| WEEDS CONTROLLED                             | WEED GROWTH STAGE               | STATE           | RATE  | CRITICAL COMMENTS   |
| Bitterbark<br>( <i>Alstonia constricta</i> ) | From summer to<br>end of autumn | NT, Qld<br>only | <p>1:4 (1 part<br/>Grazon Extra<br/>to 4 parts<br/>water)</p> <p>2 % solution<br/>for spot spray<br/>(eg. 100 mL<br/>Grazon Extra<br/>in 5 L water)</p> | <p>For use with blanket wipers only.</p> <p>For best results apply in autumn to tall (&gt; 60 cm) plants using two opposite directional passes (up and back).</p> <p>Follow up “missed” plants with a spot spray application. These will be obvious after 6 weeks.</p> <p>Blanket wiper applications can be made in summer when plants are smaller but follow up spot spraying may be necessary.</p> <p>Do not disturb (cultivate) the treated patches for at least 3 months. Best long term control is achieved when patches are left undisturbed for as long as possible after treatment (at least 6 months).</p> <p><b>Spot spraying “missed” plants:</b> thoroughly wet all stems and leaves without producing any solution run-off. Avoid any spray reaching the soil surface.</p> |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**





## IN TASMANIA FOR BLACKBERRY

**DO NOT** treat bushes carrying mature or near mature fruit.

## FOR NATIVE VEGETATION

Use of Grazon Extra on native vegetation must be done in accordance with STATE and/or LOCAL legislation.

## WITHHOLDING PERIOD

### Pasture:

Cutting or Grazing Pastures for Stock Food: **NOT REQUIRED WHEN USED AS DIRECTED.**

### LIVESTOCK DESTINED FOR EXPORT MARKETS

The grazing withholding period only applies to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that in addition to complying with the grazing withholding period, the Export Slaughter Interval is observed before stock are sold or slaughtered.

### EXPORT SLAUGHTER INTERVAL (ESI) – 3 days:

Livestock that has been grazed on or fed treated crops within 42 days of application should be placed on clean feed for 3 days prior to slaughter.

### EXPORT GRAZING INTERVAL (EGI) – 42 days.

Livestock that has been grazing on treated crops or pasture should not be sold for export slaughter for 42 days (6 weeks) after application of the chemical product, unless the export slaughter interval has been observed. When Grazon Extra is used as directed and the above withholding periods and/or export intervals are observed, treated grain and livestock commodities are considered acceptable for export. However, export requirements are subject to change. Consult your exporter for updated information about specific market requirements.

**IMPORTANT: Read the MANAGEMENT OF RESIDUES IN COMPOST, MULCHES AND ANIMAL WASTE in the PROTECTION OF CROPS, NATIVES AND OTHER NON-TARGET PLANTS section of this label.**

## GENERAL INSTRUCTIONS

### MINIMUM RECROPPING PERIODS

Aminopyralid and picloram remain active in the soil for extended periods depending on the rate of application, soil type (clay content), rainfall, temperature, humidity, soil moisture and soil organic matter. The following tables show plant-back periods to particular crops following application of Grazon Extra in different areas/situations of Australia.

**Table A: Northern New South Wales & Queensland**

Plant-back periods for rotational crops following application of Grazon Extra for rates up to 600 mL/ha on black cracking clay soils. These plant-back periods are based on a normal rainfall pattern. During drought conditions (or when the rainfall is less than 100 mm for a period of 4 months or greater) the plant-back period may be significantly longer.

| Plant-back periods for crops following the application of Grazon Extra for rates up to 600 mL/ha. |        |     |     |     |
|---|--------|-----|-----|-----|
| RATE mL/ha  | 200    | 300 | 400 | 600 |
| CROP  | Months |     |     |     |
| Wheat   | 2      | 2   | 4   | 4   |
| Barley  | 2      | 2   | 4   | 4   |
| Canola  | 2      | 4   | 4   | 4   |
| Faba bean   | 4      | 4   | 6   | 6   |
| Chickpea  | 4      | 6   | 6   | 6   |
| Lucerne   | 6      | 9   | 9   | 9   |



**Table B: Blanket Wiper Application**

| Plant-back periods for crops following blanket wiper application. |   |
|---|---|
| CROP  | Months  |
| Broadleaf crops   | 18  |
| Lucerne   | 6<br>This will allow any potential soil residues to dissipate, if any, and allow effective control. |

**Table C: Southern New South Wales**

Plant-back periods for rotational crops following application of Grazon Extra for rates up to 500 mL/ha.

| CROP   | Plant-back Period (months) |
|--|----------------------------|
| Barley, Canola, Wheat  | 9                          |
| Chickpea, Faba bean, Field pea, Lucerne, Lupin, Medic, Subclover | 24                         |

**Note:** Before using Grazon Extra in tank mixes with other herbicides, check the plant-back information on all product labels. The most residual product, i.e. the product with the longest plant-back period, will determine the time between spraying and planting.

## COMPATIBILITY

Follow any regional restrictions, and all directions and restrictions on the label, of any chemical mixed with Grazon Extra (e.g. 2,4,-D amine).

Grazon Extra is compatible with the following **herbicides**: Dow AgroSciences Amine 625,(2,4-D), Abound™ 400, AgriCrop Lantana 600, Ripper™ 480, glyphosate, metsulfuron-methyl, Roundup® CT, Starane™ Advanced.

Grazon Extra is compatible with the following **adjuvants**, as per Directions for Use on this label; Uptake™ Spraying Oil, BS1000 Biodegradable Surfactant, (alternatives are Chemwet 1000, and Spreadwet 1000 Wetting Agent), or Pulse® Penetrant.

Not all surfactants or crop oils are of equal quality, Dow AgroSciences does not support the use of alternative products other than those listed in the COMPATIBILITY section.

## MIXING

### Mix only with water.

Half fill the spray unit with water, and add the required amount of Grazon Extra. Add the remaining water with the agitator running. If required, then add spray oils or wetters (surfactants). Maintain mechanical or by-pass agitation in the spray tank during spraying. Only mix sufficient solution for immediate daily use and avoid storing.

## APPLICATION

**Only apply Grazon Extra under atmospheric conditions that do not allow drift onto sensitive crops to occur.**

### 1. WOODY WEED SITUATIONS

**Weeds need to be actively growing for herbicides to have optimum effect. Delay treatment until all regrowth has had time to grow to approximately 1 metre in situations which have been bulldozed, slashed, burnt, ploughed or areas having a previous chemical treatment.**

#### A. High Volume Spraying

Thorough coverage of foliage to the point of run-off is essential; however, avoid excess spraying which is wasteful of chemical.

#### Hand Gun

Apply the recommended mix to give full coverage of leaves and stems through a No. 6 to 8 tip at 700 to 1500 kPa (400 to 500 kPa for St John's wort). A spray volume of 3000 to 4000 L per infested hectare of 1 to 2 metre high blackberry (30 to 40 L/100 m<sup>2</sup>) should be used.

Use 2000 L of spray mixture/ha of galenia infestation (i.e. 20 L /100 m<sup>2</sup> infested area).

#### Knapsack & 12 volt Sprayer Packs

Apply the recommended spray mix to give full coverage of leaves and stems.

Only recommended for the control of herbaceous weeds, such as capeweed, fireweed and spear thistle, and woody weeds that are not regrowth less than 60 cm high or 60 cm diameter.



## B. Aerial Application

Apply in 200 L of water/ha using an aircraft to apply 100 L per pass on a double overlap pattern using nozzle configurations to produce coarse to very coarse droplets as defined by ASABE S572.

## C. Controlled Droplet Application (C.D.A.)

Results similar to high volume spraying can be obtained using Micron Herbi® or similar equipment. Select a nozzle to give a flow rate of 2 mL/sec and sweeping action of approximately 1 m/sec to ensure a droplet density of 20/cm<sup>2</sup>. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage. Also, consult directions provided with C.D.A. unit.

## D. Low Volume High Concentrate Application Techniques

Good control will be achieved, similar to high volume application, where bush size enables good coverage of entire bush. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage.

**Gas powered gun:** Apply 50 mL shots to obtain uniform coverage of 4 to 5 m<sup>2</sup> of surface area of bush. This relates to 20 droplets/cm<sup>2</sup> of leaf surface.

**Sprinkler sprayer:** This technique involves using a micro sprinkler that is connected to a hollow fibre glass rod attached to a pneumatic knapsack sprayer. Use at low pressures (50 to 200 kPa) and apply with a slow sweeping action over the top of the plants ensuring even coverage on the leaves.

## E. Boom Application

Application in a minimum spray volume of 200 L/ha for galenia and St. John's wort and 600 L of water/ha for sickle pod. Use nozzle configurations to produce medium droplets as defined by ASABE S572. Boom height must be set to ensure double overlap of nozzle patterns.

## 2. FALLOW SITUATIONS

### A. Boom Application

Application of Grazon Extra in a minimum spray volume of 50 L/ha is recommended using nozzle configurations to produce medium droplets as defined by ASABE S572. Boom height must be set to ensure double overlap of nozzle patterns.

### B. Blanket Wiper Application

Blanket needs to be made from durable and wettable material with a rigid backing. Blanket should be rigidly mounted behind motorised vehicle (eg. tractor, 4-wheel drive vehicle) and set low but never touching the ground. The chemical solution should be fed to the blanket at a flow rate sufficient to keep the blanket wet but not dripping. In thick patches

the blanket may require more frequent solution recharge (rewetting).

Ideally, a scraper bar should be mounted in front of the blanket in order to scrape or damage the bark (but not sever the stems) prior to the blanket wiping the stems. This scraper may be mounted at the front of the vehicle. Two passes (in opposite direction) with the blanket increases the contact with the plant. Ground speeds of 10-15 kph are ideal for blanket wiping application.

## CLEANING SPRAY EQUIPMENT

### Rinsing

After using Grazon Extra, empty the spray unit completely and drain the whole system. Thoroughly wash inside the unit using a pressure hose. Drain the spray unit, and clean any filters in the tank, pump, lines, hoses and nozzles.

After cleaning the spray unit as above, quarter fill with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

### Decontamination

Before spraying cotton and other sensitive crops with equipment that has been used to apply Grazon Extra (see PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section).

Wash the tank and rinse the system as above. Then quarter fill the tank and add a standard alkali based laundry detergent at 500 g (or mL)/100 L water and circulate throughout the system for at least 15 minutes. If using a concentrated laundry detergent use 250 g (or mL)/100 L water. Do not use chlorine based cleaners.

Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

**Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.**

## RESISTANT WEEDS WARNING

### GROUP I HERBICIDE

Grazon Extra Herbicide is a member of the pyridines group of herbicides. The product has the disrupters of plant cell growth mode of action. For weed resistance management, the product is a Group I herbicide. Some naturally-occurring weed biotypes resistant to the product and other Group I herbicides may exist through



normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by the product or other Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Dow AgroSciences accepts no liability for any losses that may result from the failure of the product to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local Dow AgroSciences representative.

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Crops susceptible to Grazon Extra include, but are not limited to: peas, lupins, lucerne, navy beans, peanuts, soybeans and other legumes, cotton, flowers, fruit, hops, ornamentals, shade trees and *Pinus* spp., potatoes, safflower, sugar beet, sunflowers, tobacco, tomatoes, vegetables and vines.

Grazon Extra is damaging to susceptible crops during both growing and dormant periods. Grasses are normally unaffected and establish quickly after treatment. Picloram and aminopyralid, two of the active constituents in this product can remain active in the soil for extended periods depending on soil type and application rate, rainfall, temperature, humidity, soil moisture and soil organic matter.

**DO NOT** apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands, pastures, waterways or native vegetation.

**DO NOT** allow physical spray drift onto waterways, native vegetation and susceptible crops.

**DO NOT** apply close to, or in areas, containing roots of desirable vegetation, where treated soil may be washed onto areas growing (or areas to be planted with) desirable plants.

**DO NOT** apply on sites where surface water from heavy rain can be expected to run off to areas containing, or to be planted with susceptible crops or plants.

**DO NOT** move soil, which may have been treated to areas where desirable plants are to be grown.

## MANAGEMENT OF RESIDUES IN COMPOST, MULCHES AND ANIMAL WASTE

**DO NOT** cut pasture for hay or silage production within 6 months of application, where it is intended for use off-farm.

**DO NOT** use treated plant material to make, compost, mulches or mushroom substrate.

**DO NOT** send straw from treated pastures off-farm for these purposes or for animal bedding.

**DO NOT** send animal manure, dairy shed and feed pad effluent that has been collected from animals grazing or fed pastures treated within the previous 6 months off-farm. Spreading/irrigating this manure/effluent may cause damage to clover and other susceptible plants.

**DO NOT** send compost made from animal waste that has been collected from animals grazing or fed pastures treated within the previous 6 months off-farm. Such compost may cause damage to clover and other susceptible plants.

**DO NOT** apply animal waste (e.g. manure, slurry) collected from animals grazing or fed pastures treated within the previous 6 months to susceptible plants or land to be used to grow susceptible plants.

**DO NOT** grow susceptible plants within the relevant plantback period in fields treated with manure/effluent from farms where animals have grazed or been fed treated pasture until a field bioassay shows there are no residues in the soil at levels injurious to the susceptible plants (see the crop rotation section).

To promote herbicide decomposition, manure should be evenly incorporated in the surface soil. Breakdown of residues in decomposing plants or manure is more rapid under warm, moist soil conditions and may be enhanced by supplemental irrigation.

## SOIL BIOASSAY

A simple bioassay can be conducted by collecting at least 10 spade spits of soil to a depth of 200 mm from around the paddock and thoroughly mixing the soil together. Place some of this soil in a shallow container to a depth of 3-5 cm and sow 100 seeds of the susceptible plant to be grown (subterranean or white clover is a good indicator plant where it is not practical to use the susceptible crop) into the soil. Keep in a warm and well light location and ensure the soil does not dry out. After plant emergence, check the number of plants that have germinated and seedling vigour. Symptoms of Grazon residues include non-germination or low plant emergence, leaf cupping, leaf whitening, stem elongation and twisting. If these symptoms occur do not grow the susceptible plant. Repeat the bioassay again after a further time interval.

Further information on residues in composts, mulches and animal wastes can be found at [www.dowagrosciences.com.au](http://www.dowagrosciences.com.au)



## PROTECTION OF LIVESTOCK

Poisonous plants may become more palatable after spraying and stock should be kept away from these plants until they have died down.

Many plants remain poisonous after death, and stock should not be allowed access, as there is a likelihood that they may graze the dead material. Such material should be burnt if possible.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

**DO NOT** contaminate streams, rivers or waterways with the chemical or used containers.

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

**DO NOT** store near food, feedstuffs, fertilisers or seed.

The method of disposal of the container depends on the container type. Read the STORAGE AND DISPOSAL instructions on the label that is attached to the container.

## SPILL AND LEAK MANAGEMENT

Do not touch or walk through spilled material. Wear a face shield or goggles, overalls buttoned to neck and wrist, chemical resistant gloves and footwear. Stop leak when safe to do so. Dam area and prevent entry into waterways, and drains.

**Small spills/leaks:** Absorb with material such as sand, soil or sawdust. Collect spilled product and place in sealable container for disposal. Spill residues may be cleaned using water and detergent. Contain and absorb wash water for disposal. Absorb and collect washings and place in the same sealable container for disposal. Dam the area of large spills and report them to Dow AgroSciences Emergency Services at 1-800 033 882.

## SAFETY DIRECTIONS

- Will irritate the eyes and skin.
- Avoid contact with the eyes and skin.
- If product in eyes, wash it out immediately with water.
- When opening the container and preparing the spray, wear cotton overalls buttoned to the neck and wrists, a washable hat, elbow-length PVC gloves and a face shield or goggles. When using the prepared spray, wear cotton overalls buttoned to the neck and wrists, a washable hat, elbow-length PVC gloves and a face shield or goggles.
- After each day's use, wash gloves, face shield or goggles and contaminated clothing.
- Wash hands after use.

## FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone: *Australia* 13 11 26.

If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

## MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet for **GRAZON™ EXTRA HERBICIDE** which is available from Dow AgroSciences on request.

Call Customer Service Toll Free on 1-800 700 096 or visit [www.dowagrosciences.com.au](http://www.dowagrosciences.com.au)

® <sup>TM</sup> Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

APVMA Approval No: 60830/53997

**EMERGENCY RESPONSE  
(ALL HOURS)**  
RING FROM ANYWHERE IN  
AUSTRALIA  
**1800 033 882**  
(LOCAL CALL FEE ONLY)

IN A TRANSPORT  
EMERGENCY ONLY  
**DIAL 000**  
FOR POLICE OR  
FIRE BRIGADE



Dow AgroSciences